



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

An Enumeration of the Plants Collected by Dr. H. H. Rusby in South America, 1885-1886.—XVI.

(Continued from p. 38.)

- MANETTIA (?) DIFFUSA, sp. nov. Ramulis gracillimis glabris, teretibus; foliis gracile petiolatis, ovatis tenuis glabris acutis vel acuminatis, 5-6 cm. longis, basi obtusis vel cordatis; inflorescentia axillari et terminali, cymosa-paniculata; pedunculis pedicellique gracilibus, glabris; bracteis minutis; corolla 6mm. longa, cylindræa, tubus basi constricta. Resembling *M. paniculata*, Poepp. and Endl. in habit but very different from that plant. Guanai, 2,000 ft. (2121).
- Cosmibuena obtusifolia*, R. & P. Fl. Peruv. iii. 3. Mapiri, 2,500 ft. (2103).
- Condaminea corymbosa* (R. & P.), D.C. Prodr. iv. 402. Yungas, 6,000 ft. (1898).
- Chimarrhis*, sp. Yungas, 6,000 ft. (2446).
- Chimarrhis*, sp. Same locality (2447).
- Pogonopus tubulosus* (D.C.), Schum. in Mart. Fl. Bras. vi. Pars. vi. (2), 265. Guanai, 2,000 ft. (2090).
- Warszewiczia coccinea* (D.C.), Klotzch, Mon. Ber. Akad. Wiss. Berl. 1853, 496. Guanai, 2,000 ft. (1953); Yungas, 6,000 ft. (1954); Falls of the Madeira, Brazil (1955).
- Sipanea pratensis*, Aubl. Guian. i. 147. (*S. hispida*, Benth. Hb. Kew). Mapiri, 2,500 ft. (2461).
- Oldenlandia herbacea*, D.C. Prodr. iv. 425. Falls of the Madeira, Brazil (2156).
- Isertia bullata*, Schum. loc. cit. 286 (?). Mapiri, 2,500 ft. (1895).  
I have not seen the type of Herr Schumann's species described as having cordate leaves. The leaves of our plant are narrowed at the base, otherwise much like those described by him.
- Sabicea cana*, Hook. f. Ic. Pl. t. 247. Mapiri, 2,500 ft. (1905).
- Sabicea aspera*, Aubl. Guian. 194, t. 76. (*S. hirsuta*, H.B.K.) Guanai, 2,000 ft. (1897); Mapiri, 2,500 ft. (1904).
- Coccocypselum canescens*, Willd. ex Cham. & Schlecht. Linnæa, iv. 139. Yungas, 4,000 ft. (1397); Mapiri, 2,500 ft. (1896); Falls of the Madeira, Brazil (2099).
- COCCOCYPSELUM MACROPODUM (R. & P.) (*Psychotria macropoda*, (R. & P.) Mapiri, 2,500 ft. (2098).

COCCOCYPSELUM GLABRUM. sp. nov. Herba ascendens, caulibus acutangulis glabris, 15-20 cm. altus. Petiolus 1 cm. longus; stipulæ subulatæ; folia ovato-lanceolata, acuta, basi obtusa, 4-6 cm. longa, 2-3 cm. lata, glabra; inflorescentia capitata, 5-10 mm. diametro; pedunculus 2 cm. longus; bacca 2 mm. diametro.

Yungas, 4,000 ft. (2479). The same as Matthew's Nos. 850 and 1943 from Peru, Herb. Kew.

*Hamelia patens*, Jacq. Stirp. Amer. 74, t. 50. Mapiri, 5,000 ft. (2105).

HOFFMANNIA BRACHYCARPA, sp. nova. Ramis adscendentibus vel erectus obtusangulis; foliis breve petiolatis vel sessilibus, obovatis, apice acuminatis basi cuneatis, supra glabris, subtus minute ferrugineo-pubescentibus, 20-30 cm. longis, 7-8 cm. longis, margine integris; inflorescentia axillaris, pedunculi gracili, 2 cm. longi; corolla ovarium aequale; bacca 2½ mm. longa 2 mm. lata. Yungas, 4,000 ft. (2522).

*Bertiera Guianensis*, Aubl. Guian. i. 180, t. 69. Guanai, 2,000 ft. (2118).

*Posoqueria longiflora*, Aubl. Guian. i. 134. t. 51. Junction of the Rivers Beni and Madre de Dios (2220).

*Basanacantha spinosa* (Jacq.), Schum. in Mart. Fl. Bras. vi. Pars. vi. 2, 376. Junction of the Rivers Beni and Madre de Dios (2157) (*Randia glabrescens*, Spruce).

*Genipa Americana*, L. Sp. Pl. Ed. 2, 251, var. *latifolia*, Spruce, Herb. Kew. Junction of the Rivers Beni and Madre de Dios (1899).

*Chomelia paniculata*, Benth. in Herb. Kew. Junction of the Rivers Beni and Madre de Dios (1878); Mapiri, 2,500 ft. (2117).

*Chomelia*, sp. Falls of the Madeira, Brazil (1490). I determined this at Kew as "*C. pubescens*, Benth" but there is a *C. pubescens*, C. & S. described in Schumann's Rubiaceæ of the Brazil Flora, p. 36, with which the plant does not agree.

*Chomelia*, sp. Guanai, 2,000 ft. (2659).

*Coffea Arabica*, L. Unduavi, 8,000 ft. Cultivated. (1900).

*Coussarea*, apparently undescribed. Junction of the Rivers Beni and Madre de Dios (2402).

*Coussarea*, sp. Beni River (2677) collected in fruit only.

- Coussarea?* Mapiri, 2,500 ft. (2559). The same as Fendler's No. 1990, from Tovar, Venezuela.
- Faramaea salicifolia*, Presl. Symb. Bot. 24, t. 70. Yungas, 6,000 ft. (1869); Unduavi, 8,000 ft. (1870); Mapiri 2,500 ft. (2120), (2626), the later collected in fruit and referred to this species with some hesitation.
- Faramaea Montevidensis*, D.C. Prodr. iv. 497. Junction of the Rivers Beni and Madre de Dios (1874).
- Faramaea brevisflora*, Benth. in Herb. Kew. Falls of Madeira, Brazil. (2571.) Name not in Schumann's Rubiaceæ of the Brazil Flora.
- Psychotria alba*, R. & P. Fl. Peruv. ii. 58, t. 205. (*Mapouria alba*, Muell. Arg.) Yungas, 4,000 ft. (2110) (1866); Guanai, 2,000 ft. (1889); Junction of Rivers Beni and Madre de Dios (1877) (1875).
- Psychotria Casiquiaria*, Muell. Arg. in Schum. Mart. Fl. Bras. vi. Pars v. 324. Mapiri, 2,500 ft. (2109). The same as Spruce, Rio Negro, No. 3436.
- Psychotria lupulina*, Benth. in Hook. Journ. Bot. iii. 230. Falls of the Madeira, Brazil (1873); Beni River (1872).
- Psychotria brachybotra*, Muell. Arg. in Schum. loc. cit. 327. Yungas, 6,000 ft (1887). The same as Spruce, Rio Negro, No. 2190.
- Psychotria barbiflora*, D.C. Prodr. iv. 509? Mapiri, 2,500 ft. (2112). Specimens too old for certain determination.
- Psychotria Marcgravii*, Spreng. Syst. Cur. Post. 79. Yungas, 6,000 ft. (1893).
- Psychotria Paraensis*. Muell. Arg. in Schum. loc. cit. 244. Falls of Madeira, Brazil (2114).
- Psychotria brachyloba*, Muell. Arg. loc. cit.? Mapiri, 5,000 ft. (1864).
- PSYCHOTRIA CHIONANTHA (D.C.) (*Palicourea chionantha*, D.C. Prodr. iv. 526; *Psychotria Luschnathii*, Mart. Herb. Fl. Bras. 311). Mapiri, 5,000 ft. (1883).
- PSYCHOTRIA BRACHYPODA (Muell. Arg.) (*Mapouria brachypoda*, Muell. Arg. in Schum. loc. cit. 422). Mapiri 2,500 ft. (1882). The same as Burchell's No. 3318 from San Paulo, Brazil.

- Psychotria flexuosa*, Willd. Sp. Pl. i. 966. Mapiri, 5,000 ft. (1884).  
*Psychotria racemosa*, Willd. Sp. Pl. i. 966. Mapiri, 2,500 ft. (1867).  
*Psychotria tabacifolia*, Muell. Arg. loc. cit. 236. Mapiri, 2,500 ft. (1880).  
*Psychotria viridis*, R. & P. Fl. Per. t. 210. Guanai, 2,000 ft. (2515). The same as Matthews, No. 1949, Peru, Herb. Kew.  
 PSYCHOTRIA CROCEA (Schlecht.) *Palicourea crocea*, Schlecht, Linnæa, xxviii. 525 Beni River (2116).  
*Psychotria pilosiuscula*, Griseb. Mapiri, 2,500 ft. (2480).  
*Psychotria crassa*, Benth. Mapiri, 2,500 ft. (2113).  
*Psychotria cornigera*, Benth. Yungas, 4,000 ft. (1865; 1881).  
*Psychotria*, near *P. venulosa*, Muell. Arg. Falls of Madeira, Brazil (1879).  
*Psychotria* near *P. idotricha*, Muell. Arg. Mapiri, 2,500 ft. (2489).  
*Psychotria leiocarpa*, C. & S. Linnæa, 1829, 22. Guanai, 2,000 ft. (2161).  
 PSYCHOTRIA NIVEO-BARBATA (Muell. Arg.) (*Mapouria niveo-barbata*, Muell. Arg.; Schum. in Mart. Fl. Bras. vi. Pars. vi. 2. 401). Mapiri, 2,500 ft. (1886). The same as Glaziou's No. 7684 from Rio Janeiro, Herb. Kew.  
 PSYCHOTRIA UMBROSA (Muell. Arg.) (*Mapouria umbrosa*, Muell. Arg. Flora 1876, 459). Mapiri, 2,500 ft. (1868). The same as Spruce's No. 660, Herb. Kew.  
*Psychotria*. Guanai, 2,000 ft. (2558).  
*Psychotria*. Mapiri, 2,500 ft. (1894).  
*Psychotria* (?) Yungas, 6,000 ft. (1885).  
*Psychotria*. Junction of the Rivers Beni and Madre de Dios (2575).  
*Psychotria*. Mapiri, 2,500 ft. (1871).  
*Psychotria tomentosa* (Willd.), Muell. Arg., Schum. loc. cit. 370. Mapiri, 2,500 ft. (853); Mapiri, 5,000 ft. (854; 855).  
*Rudgea micrantha*, Muell. Arg. Flora, 1876, 454. Falls of the Madeira, Brazil (2108).  
*Rudgea Amazonica*, Muell. Arg. loc. cit. 449. Junction of the Rivers Beni and Madre de Dios (1890).  
*Rudgea Hostmanniana*, Benth. Yungas, 4,000 ft. (1888; 1892). Mapiri, 5,000 ft. (1891).  
*Rudgea*. Mapiri, 2,500 ft. (1876).